

# Increased circularity and reduced CO<sub>2</sub> footprint of concrete production by use of industrial wastes, calcinated clays and bio-ashes

Katarina Malaga, prof. dr.  
Vice President, RISE Research Institute of  
Sweden  
Head of CBI Swedish Cement and Concrete  
Institute

2019-12-03, Brussels

# Urbanisation and increased demand of cement

- Cement production: 4800 Mt 2018 - 6% of the total anthropogenic CO<sub>2</sub> emissions
- Concrete production: 30000 Mt 2015
- Urbanisation and economic development increase
- 2050 - 3 mld people from poor areas in need of housing






Figure 2: Cement production and emissions, 2010–15




From: A sustainable future for the European cement and concrete industry; ETH/EPFL, 2018

# Embodied emissions and energy for materials used in construction (UK)



From: A sustainable future for the European cement and concrete industry; ETH/EPFL, 2018

# How can we reduce the CO<sub>2</sub> emissions in concrete production?



# Less cement – less natural resources - lower CO<sub>2</sub>

- CDW – construction demolition waste
- SCMs – supplementary cementitious materials
  - fly-ash, slag, bio-ashes, clays

# Use of CDW in concrete

## Some objectives

- Reuse of CDW in real construction
- Lower use of natural resources
- Development of prefab elements integrating a high level of CDW-derived materials
- To build a house of CDW

<http://www.re4.eu/>

### REUSABLE MATERIALS






### CONSTRUCTION & DEMOLITION WASTE



### RE<sup>4</sup> SORTING SYSTEM




PREFABRICATION OF ELEMENTS FOR BUILDINGS  
REFURBISHMENT & CONSTRUCTION



Up to 80-90% of reusable structures for RE<sup>4</sup>-prefabricated building concept

# Sorting of CDW → valuable resources



# CDW as concrete aggregate



8/16



2/8



0/2


Recycled aggregates from the CDE recycling centre outside Marseille, southern France (fraction limits in mm)

# New CDW quality classes – a proposal

## Instead 2 classes → 4

Property	Type A+	Type A	Type B	Type L
Density (oven dry) [kg/m <sup>3</sup> ]	> 2000	> 2000	> 2000	≤ 2000 (To be declared)
Rc+Ru [%]	≥ 95 %	≥ 90 %	≥ 70 %	---
Rb [%]	≤ 5 %	≤ 10 %	≤ 30	---
Rc+Ru+Rb [%]	---	----	≥ 90 %	≥ 95 %
Ra [%]	≤ 1 %	≤ 5 %	≤ 10 %	≤ 5 %
FL	≤ 0,2 %	≤ 2 %	≤ 2 %	≤ 5 %
X+Rg [%]	≤ 1 %	≤ 5 %	≤ 5 %	≤ 5 %
Shape	≤ Fl <sub>35</sub> or ≤ Sl <sub>40</sub>	≤ Fl <sub>50</sub> or ≤ Sl <sub>55</sub>	To be declared	To be declared
Water soluble sulfates	≤ SS <sub>0,2</sub>	≤ SS <sub>0,2</sub>	≤ SS <sub>0,2</sub>	≤ SS <sub>0,2</sub>
Influence on setting time	≤ A <sub>10</sub>	≤ A <sub>40</sub>	≤ A <sub>40</sub>	≤ A <sub>40</sub>
Water absorption	≤ 5%	≤ 10 %	To be declared	To be delared


# Sandwich element production – upscaling at Creagh Concrete



- Self-Compacting Concrete with 50% of aggregates being CDW (Type B)
- High-Performance Concrete with 50% of aggregates being CDW (Type B)
- With CDW quality class Type A, usage levels were 80 and 90 %, respectively



# Achievements: Prefab elements constituting 30-95 % CDW



# Demo buildings in Spain and UK (below from Creagh Concrete, UK)



Use of clays and bio  
ashes in concrete:

The path towards a  
better sustainability and  
increased circular  
economy



# Which SCM?


- Chemical composition
- Availability
- Reactivity

- Commonly use SCMs are waste products from other industries:
  - they are often imported
  - have a limited availability
  - shortage is forecasted

- New sources of SCMs are needed, ideally local ones

→ Clays

→ Bioashes




# Swedish clays as SCM

- Availability ✓
- Composition ✓
- Reactivity

Available in many part of the world, including Sweden

-  **West Coast**  
Marine glacial and postglacial clays
-  **Eastern central Sweden**  
Brackish marine glacial clays, freshwater deposited clays occur
-  **Littorina Sea**  
Brackish marine postglacial clays, including sulfide clays
-  Areas below the highest shoreline, where the salinity is expected to have been at its lowest during clay deposition
-  Areas above the highest shoreline



SiO<sub>2</sub> Their chemical composition make them compatible with cement



Al<sub>2</sub>O<sub>3</sub>

CaO

# Clay as SCM

- Availability ✓
- Composition ✓
- Reactivity

- To be used with cement, clays must be activated, usually by heating at 600-800°C




# Clay as SCM

- Can be used together with limestone to reduce even more the amount of cement (up to 50% replacement)
- Swedish clays show promising results: good reactivity and mechanical strength

- Availability ✓
- Composition ✓
- Reactivity ✓

Compressive strength



# Bio-ashes as SCM

- Availability ✓
- Composition
- Reactivity

- According to Energiforsk:  
1.7 M t/ year of ashes is produced by Swedish energy production  
Bottom ash – 0.7 M t a year has a priority to be used in other industries
- Pulp and paper industry: approx. 1.5 M t/year,  
most of it is landfilled.


Can we use this waste to replace cement,  
and contribute to a circular economy?



# Bio ashes as SCM

- Availability ✓
- Composition ✓
- Reactivity

- Chemical requirements:
  - The chemical composition should not vary over time to ensure a constant quality
  - No chloride (washable if present), no heavy metals




→ Check and approved for 7 different Swedish companies, with sampling up to 6 months

# Bio-ashes as SCM


- Chemical composition ✓
- Availability ✓
- Reactivity ✓

- The tested bio-ashes exhibit good reactivity with cement

Example fly ash




Example bottom ash



→ Development of binders in process

# A possible future use of cement?



**Thank you for your attention!**

**[katarina.malaga@ri.se](mailto:katarina.malaga@ri.se)**

